

# PropiMax\* EC

## Section 18 Quarantine Exemption

### For Control of Soybean Rust (*Phakopsora pachyrhizi*) in Soybean

#### For Distribution and Use Only in the State of Mississippi

#### ATTENTION

- **This quarantine exemption expires March 1, 2007.**
- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- The product PropiMax\* EC fungicide, EPA Reg. No. 62719-346, (containing 41.8% propiconazole) manufactured by Dow AgroSciences LLC may be used.
- All applicable directions, restrictions and precautions on the Section 18 use directions as well as the EPA-registered product label for PropiMax EC must be followed.

#### Directions for Use

Apply PropiMax EC at the rate of 4 - 8 fl oz per acre. PropiMax EC may be applied using ground or aerial broadcast equipment or through sprinkler irrigation (chemigation) equipment. Use sufficient spray volume to provide for uniform coverage and canopy penetration, normally 15 to 20 gallons per acre (gpa) by ground or 5 or more gpa by air. Best results are obtained when PropiMax EC is applied at the first sign of disease. If the disease is established, use the higher rate within the rate range. A single repeat application may be made at an interval of 10 to 14 days after the first.

**Chemigation:** Apply PropiMax EC in 1/4 to 1/2 inches of sprinkler irrigation. Follow recommendations for application rates, timing and frequency of application in crop-specific directions for use (above).

#### Restrictions:

- Use of PropiMax EC under this quarantine exemption will not be allowed until the time that soybean rust has been confirmed to be present in the continental United States by the U.S. Department of Agriculture-Animal and Plant Health Inspection Service (USDA/APHIS). Once USDA/APHIS has confirmed that soybean rust is present, all existing EPA-approved section 18s for soybean rust will be automatically released for implementation in Mississippi as the Mississippi Department of Agriculture & Commerce feels is appropriate.
- **Do not** make more than 2 applications or apply more than 12 ounces (0.338 lb active ingredient) per acre per year.
- Do not apply after R5 (pod fill) stage of growth.
- **Do not** use hay, forage, or fodder from the soybean crop as any component of animal feed or bedding.
- **Rotational Crop Restrictions:** To avoid possible illegal residues, do not plant any rotational crop intended for food, grazing, or any component of animal feed or bedding within 105 days of application of PropiMax EC to the preceding crop, unless the rotational crop appears on the label for PropiMax EC.
- A maximum of 2 total applications using approved Section 18 products collectively are allowed under this soybean rust Section 18 exemption.

Any adverse effects resulting from the use of PropiMax EC under this emergency exemption must be immediately reported to the Mississippi Department of Agriculture & Commerce.

\*Trademark of Dow AgroSciences LLC

R-142-004

EPA-accepted: 07-27-2004

File symbol: 04-MS-17

Initial printing.